

THE EFFECT OF CYANOBACTERIAL TOXIN ON SEED GERMINATION

R.M.S. SENGAR^{a1}, SEEMA BHADAURIA^b AND POONAM SHARMA^c

^aDepartment of Botany, Agra College, Agra, U.P., India
E-mail: rudra_sengar@rediffmail.com

^bDepartment of Botany, R. B. S. College, Agra, U.P., India
E-mail: bhadauriaseema1@rediffmail.com

^cMicrobiology Research Lab, Department of Botany, Agra College, Agra, U.P., India

ABSTRACT

The world-wide occurrence of harmful cyanobacterial blooms CyanoHAB in fresh and brackish waters create problems for all life forms. During CyanoHAB events, toxic cyanobacteria produce cyanotoxins at high level under stress conditions that can cause chronic and sub-chronic toxicities to animals, plants and humans. Cyanotoxicity in eukaryotes has been mainly focused on animals, but during recent years, data, related to cyanotoxin impact on both aquatic and terrestrial crop plants irrigated by water containing these toxins have become more and more available. This last cited fact is gaining importance since plants could in a direct or indirect manner contribute to cyanotoxin transfer through the food chain, and thus constitute a potent health risk source. The use of this contaminated irrigation water can also have an economical impact which appears by a reduction of the germination rate of seeds, and alteration of the quality and productivity of crop plants. The main objective of present investigations is to discuss the eventual phytotoxicity of cyanotoxin as the major agricultural impacts induced by the use of contaminated water for plant irrigation. Thus cyanobacterial phytotoxicity strongly suggests a need for the surveillance of CyanoHAB and the monitoring of water irrigation quality. Therefore it is advisable to farmers that they should not use such water bodies for irrigation without prior examination.

KEY WORDS: *Microcystis aeruginosa*, *Vigna radiata*, Cyanobacterium, CyanoHAB, cyanotoxin, water irrigation

Cyanobacteria are one of the most diverse groups of Gram-negative photosynthetic prokaryotes which frequently form cyanobacterial harmful blooms (CyanoHAB) in eutrophic water bodies. Some species of cyanobacteria are able to produce toxins which may be divided in to three main groups; hepatotoxin, neurotoxin and cyanotoxin (Codd, 2000). These toxins can cause water quality problems for fisheries, aquaculture, farming and sanitary hazards for human and animals. Therefore, survey of CyanoHAB and cyanotoxin have been carried out in several countries, avoiding sanitary risks by a safe use of cyanotoxin-contaminated water (Chorus and Bartram, 1999). However, in water irrigation, cyanotoxin surveys have not yet been considered within any water quality official monitoring programme, in spite of the possible contamination of edible plants constituting a significant indirect route of human exposure to cyanobacterial toxins (Codd, 2000). Accordingly, many studies clearly indicate that irrigation with water containing cyanotoxin can be a threat for both the quality and yield of crop plants (Crush et al. 2008, Saqrane and Odura, 2009). Ko's et al. (1995) reported for the first time that cyanotoxin inhibited the growth of mustard seedlings.

Recently, the effects of cyanotoxins on seed germination and contaminated with cyanotoxin in irrigation leads to lower field productivity. In this way several terrestrial plants were investigated using both cyanobacterial standard toxin and cyanobacterial cell-free crude extract by several workers like Saqrane et al. (2008) and Crush et al. (2008). In the present investigation the attempt has been made to study the effect of cell free extract of a toxin producing cyanobacterium *Microcystis aeruginosa* on seed germination of *Vigna radiata* because it is one the most medicinally used pulse crop of this region.

MATERIALS AND METHODS

The toxic property of *Microcystis aeruginosa* was first observed in laboratory cultures while attempting to grow seeds of *Vigna radiata*. *Microcystis aeruginosa* has been isolated in unialgal form following Bold (1942) and then purified by following Watanabe and Kiyohara (1963) method. Then following methodology have used in the present investigation:

Author's formulated culture medium

Filtered and sterilized natural water	990 ml
Soil extract-w/v	10 ml/l

¹Corresponding author

This *M. aeruginosa* was cultured in the culture room at 30°C temperature with light exposure of 3200 lux for 16 hours of 24 hours cycle

Extraction of toxic substances

Author's modified medium was taken for the culture of this cyanobacterium as it was growing luxuriantly under natural conditions and the toxin extraction was done from 30 days old culture, because on this day the cyanobacterial growth was maximum followed by senescence. The investigations for the extraction of toxic substances were carried out by adopting two methods:

A. Extraction from algal cells (Ramamurthy and Krishnamurthy, 1967)

The cyanobacterium was thoroughly washed in sterilized glass distilled water and 1 ml of the packed cell volume of the selected species of cyanobacterium from fresh sub-cultures was then crushed with 2 ml of sterilized glass distilled water. A bluish green viscous fluid was there by obtained. The toxic property was tested by following Florey et al (1949) and Waksman (1952).

B. Extraction from culture filtrate

The cyanobacterial suspension along with whole culture medium was filtered through Watman's No.1 filter paper. The filtrate was collected in beaker. This filtrate was dried in oven at 30°C ± 5°C. The dried filtrate was mixed with 1 ml distilled water and collected in glass vial. The toxic property of this extract was also tested by the methods being described in the ensuing paragraph.

Assessment of cyanobacterial toxin

Since *Microcystis aeruginosa* was predominant cyanobacterial form in many aquatic reservoirs of Agra, hence it was selected for the study. This cyanobacterium was grown in culture flask in author's modified medium for 30 days and culture extract was obtained by filtration. This extract was used for the treatment of the seeds. The extract was used as such in various dilutions and proceeded as below:

1. The moong (*Vigna radiata*) seeds were soaked for 10 hours in extract of this cyanobacterium in the test tubes
2. Then the treated seeds were transferred on the blotting paper in the petridishes
3. The observations for the seed germination were taken

after every 12 hours up to 48 hours

4. Length of radicle of the germinated seeds was measured (Allen, 1973)

RESULTS AND DISCUSSION

The *M. aeruginosa* under normal conditions do not secrete toxin hence this habitat is enjoyed by cattle of nearby area. But in April-May months cattle avoid such water reservoir habitat as they sense danger of polluted habitat. It was because of foul smell and toxin production. The various factors which could induce toxin production were studied under in laboratory conditions. When it was grown at high temperature i.e. 30°C with 3200 lux of light continuously for 16 hrs of 24 hrs cycle and supply of 1% NaHCO₃ only then toxin production was there. Therefore, it became clear that stress conditions under natural environment induce toxin production. In April-May months the air temperature of Agra goes high i.e. 37-40°C which caused heating of surface water and high light condition prevail with increased concentration of salts due high evaporation. These stress factors induce *Microcystis aeruginosa* to secrete toxins. This in conformity with the results of Gorham (1960) but in contrast of Leferve (1964). Opinions vary greatly for the influence of agitation as well as CO₂ supply in the production of toxin by blooms. In the present studies daily shaking of culture flasks at (8 hrs intervals) and supply of CO₂ in the form of NaHCO₃ (1%) could give better results.

To study the effect of bloom toxicity upon seed germination the seeds of a pulse *Vigna radiata* has been selected, it was because, in case of water scarcity, farmers used pond water to irrigate their agricultural fields. After going through the observation as shown in table no-1 it became clear that with the increase of cyanobacterial cell free extract concentration the growth of radicle was found decreasing. However in higher concentration, few seeds could not germinate. These results showed the inhibitory effect of cyanotoxin upon seed germination (plate 1-2). The cell free cyanobacterial extract in different concentrations affected the germination of seed. During the course of study it was found that in the presence of cyanobacterial toxin

SENGAR ET AL. : THE EFFECT OF CYANOBACTERIAL TOXIN ON SEED GERMINATION

seed germination was found delayed and further the length of radicle decreased as the concentration of cyanobacterial extract increased. Thus the effect of cyanotoxin on seed germination was dose dependent. As the cell free extract concentration increased beyond 6ml in 4ml distil water,

the percentage of germination started decreasing. Further, it was noted that with the increase of cell-free extract concentration the length of radicle of germinated seed was affected. These results clearly showed the inhibitory role of cyanotoxin. It appeared that exposure to cyanobacterial

TABLE 1: EFFECT OF CYANOBACTERIAL EXTRACT ON SEED GERMINATION**

P D N O	CONCENTRATIONS	NO. OF SEEDS GERM INATED	LENGTH OF RADICLE OF GERMINATED SEEDS *(cm.)										AVE- RAGE
			1	2	3	4	5	6	7	8	9	10	
1	10 ml Distil Water(Control)	10	4.7	3.4	2.8	3.6	3.3	2.0	1.6	1.4	1.3	1.7	2.6
2	9 ml Distil Water + 1ml. cyanobacterial extract	10	3.1	3.1	3.6	4.2	2.5	1.4	1.9	2.8	1.4	1.1	2.5
3	8 ml Distil Water + 2 ml cyanobacterial extract	10	2.9	1.8	3.3	2.3	2.0	2.2	2.3	2.1	1.9	1.8	2.3
4	7 ml Distil Water + 3ml. cyanobacterial extract	10	3.4	3.2	2.3	2.8	2.1	2.0	1.3	1.7	1.4	1.7	2.2
5	6 ml Distil Water + 4ml. cyanobacterial extract	10	2.8	3.5	2.8	2.0	1.8	2.0	2.6	1.7	1.3	0.6	2.1
6	5 ml Distil Water + 5ml. cyanobacterial extract	10	2.8	2.2	2.0	1.8	2.0	1.5	2.5	2.0	0.8	1.4	1.9
7	4 ml Distil Water + 6ml. cyanobacterial extract	9	2.6	2.2	1.3	2.3	1.5	1.6	1.6	1.7	1.8	0.0	1.7
8	3 ml Distil Water + 7ml. cyanobacterial extract	9	2.5	2.4	1.3	1.0	1.6	1.3	0.8	2.5	2.5	0.0	1.6
9	2 ml Distil Water + 8ml. cyanobacterial extract	8	2.4	1.5	2.0	0.0	2.1	2.2	1.8	1.9	0.0	1.3	1.5
10	1 ml Distil Water + 9 ml cyanobacterial extract	7	2.0	0.0	2.0	1.5	1.8	2.0	0.0	1.9	1.5	0.0	1.3

Pd No. Petridish Number

*The Seeds Of *Vigna radiata*

**Test Organism (*Microcystis aeruginosa*)

SENGAR ET AL. : THE EFFECT OF CYANOBACTERIAL TOXIN ON SEED GERMINATION


aqueous extract may affect the metabolic activities of seeds during the germination process. Similar results were obtained by Chen et al. (2004) using rice and rape seeds. The inhibition of seed germination was also observed after the exposure of Alfalfa (*Medicago sativa*) seedling to cyanobacterial cell-free crude extract, by Pflumacher et al. (2006). All these investigations suggested that exposure to cyanotoxin via irrigation contaminated with toxins of cyanobacteria could pose a threat to the quality and yield of crop plants in the agricultural environment. Therefore, it is suggested that farmers should not use the CyanoHAB

containing water to irrigate their land without prior examination

ACKNOWLEDGEMENTS

Authors are highly obliged to the Principal, Agra College, Agra and RBS College, Agra for their continuous motivation and all possible help during the course of study.


Plate No. 1



Photograph showing effect of cyanotoxin on seed germination

A- Control against known toxin
B- Control against unknown toxin

Plate No. 2



Photograph showing effect of cyanotoxin on seed germination

A- Control against low concentration of cyanotoxin
B- Control against high concentration of cyanotoxin

REFERENCES

- Allen M. M.; 1973. Methods for Cyanophyceae. In: Handbook of Phycological Methods-Culture, Methods and growth measurements. Edited by J. R. Stein.(Cambridge Univ. Press), London:127-238.
- Bold H.C.; 1942. cf. The Cultivation of Algae, Venkatraman,G.S.1969 ICAR Publ. New Delhi: 69-138.
- Chen J., Song L., Dai J., Gan, N. and Liu Z.; 2004. Effect of Microcystin on the growth and the activity of superoxide dimutase and peroxidase of Rape (*Brassica napus* L) and rice (*Oryza sativa* L). *Toxicon*, **43**:393-400.
- Chorus I and Bartram J.;1999. Toxic cyanobacteria in water. A guide of their public health consequences, monitoring and management; E & FN Spon/WHO, London, UK: 416.
- Codd G.A., Metcalf J.S., Beattie K.A.;1999. Retention of *Microcystis aeruginosa* and microcystin by salad lettuce (*Lactuca sativa*) after spray irrigation with water containing cyanobacteria. *Toxicon*, **37**:1181-1185.
- Codd G.A.; 2000. Cyanobacterial toxins, the perception of water quality, and prioritisation of eutrophication control. *Ecol. Eng.*, **16**: 51-60.
- Crush J.R., Briggs L.R., Sprosen J.M. and Nichols S.N.; 2008. Effect of irrigation with lake water containing microcystis on microcystin content and growth of rye grass, clover, rape and lettuce. *Environ. Toxicol.*, **23**: 246-252.
- Florey H.W., Chain E., Heatley N. G., Jennings M.A., Sanders A.G., Abraham E.P. and Florey M.E.;1949. Antibiotics a survey of penicillin, streptomycin and other antimicrobial substances from fungi, actinomycetes, bacteria and plant, **2**, Oxford Univ.Press, New York.
- Gorham P.R. ;1960. Toxic water blooms of blue green algae. *Can. Vet. J.*, **1**: 235-245.
- Ko's P, Grozo G., Suranyi G. and Borbely G.; 1995. Simple and efficient method for isolation and measurement of Cynobacterial Hepatotoxins by plant tests (*Sinapsis alba* L.). *Anal. Biochem.*, **225**:49-53.
- Lefevre M. ;1964. Extracellular products of algae: In *Algae and Man*. Edited by D.F. Jackson (Plenum Press) New York.
- Pflugmacher S., Jung K., Lundvall L., Neumann S. and Peuthert A.; 2006. Effects of cyanobacterial toxins and cyanobacterial cell free crude extract on germination of Alfalfa (*Medicago sativa*) and induction of oxidative stress. *Environ. Toxicol. Chem.*, **25**:2381-2387.
- Ramamurthy V.D. and Krishnamurthy S.; 1967.The antibacterial properties of marine blue green alga *Trichodesmium erythraeum*(Her). *Curr. Sci.*,**19**: 524-525.
- Saqrane S., Ghazali E.L., Odura B., Bouarab L. and Vasconcelos V.; 2008. Effects of cyanobacteria producing microcystins on seed germination and seedling growth of several Agricultural plants. *J. Environ. Sci. Health B*, **43**:443-451
- Saqrane, S. and Odura, B.;2009. CyanoHAB occurrence and water irrigation cyanotoxin contamination: Ecological impacts and potential health risks.*Toxicon*, **1**: 113-122.
- Waksman, S.E.; 1952 *Soil microbiology*, Publ.Willy and Sons, Inc., New York, USA:267
- Watanabe,A. and Kiyohara, T.; 1963, cf. *Microalgae and photosynthetic bacteria*, Univ, Tokyo Press, Japan; 182-196.