



FLORISTIC DIVERSITY OF LAKH BAHOSI BIRD SANCTUARY- A TWIN LAKES WETLAND OF UTTAR PRADESH, INDIA

PARITOSH KUMAR TEWARI^a, SHUBHAM KUMAR^b, ATUL KUMAR ANAND^c AND ALKA KUMARI^{d1}

^{abcd}Department of Botany, University of Lucknow, Lucknow, Uttar Pradesh, India

ABSTRACT

The present paper explores floristic composition, diversity and nature of plant communities occurring in the Lakh Bahosi Bird Sanctuary, Uttar Pradesh, which is totally based on field survey and identification. During the preliminary field survey around 110 species have been collected from the study site in which the dominant and keystone genera are *Ficus bengalensis*, *Azadirachta indica*, *Alstonia scholaris*, *Terminalia arjuna* and *Syzygium jambolanum* etc. Besides, some aquatic species like *Azolla pinnata*, *Typha latifolia*, *Ipomoea aquatica*, *Nymphaea nouchali*, and *Pontederia crassipes* are also growing as dominant plant resources. Critical literature survey and studies indicates that the wetland diversity has been fast deteriorating due to over-exploitation of biological resources by human population of surrounding areas.

KEYWORDS: Protected areas, Kannauj District, Bird Sanctuary, Lakh Bahosi, Wetland

Wetlands with water logged soil conditions possess characteristic life forms, both plants and animals, which can tolerate long periods of water saturated environments. They are the sites of hydromorphic soils and have a wealth of biodiversity. Previous studies have identified their economic, ecological and cultural values. The ecological roles of wetlands have been recognized as to maintenance of water quality as well as quantity, hydrology, flood control, carbon cycle, climate stability, water cycle, recharge of ground water. The state of Uttar Pradesh in India is endowed with 12 significant wetlands constituting a part of the Upper Gangetic plains, which are proclaimed as bird sanctuaries.

Lakh Bahosi Sanctuary is located near Lakh Bahosi village in Kannauj District of Uttar Pradesh established in 1989. It is one of largest bird sanctuaries in India spread in areas of 80 sq. km. The sanctuary is a famous spot for the birdwatchers and nature enthusiasts. This area has been covered by huge numbers of flora and fauna. The wetland area has large number of aquatic plants also. Although the diversity of the area has been explored since 20 years ago by Sinha (2005) but not given much emphasis in recent years. The floristic composition of protected areas like wetlands/bird sanctuary is more significant to learn the diversity and taxonomy of the plants. Although other bird sanctuaries of Uttar Pradesh are more explored recently but this particular area has not been given more emphasis so the present study has been aimed to study its floristic composition, diversity, habit and habitat and their threat status as per IUCN categories.

MATERIALS AND METHODS

Study Site

Lakh Bahosi Wildlife Sanctuary is about 38 km from the historic city of Kannauj. The Sanctuary is formed of two oxbow Jheels near the village Bahosi. Both the Jheels, Lakh and Bahosi, are located near the Lower Ganga Canal, so the overflow and seepage of water accumulates in the Jheels, resulting in about 600 ha of shallow wetlands perfectly suitable for water birds. By winter, the water spread is reduced by evaporation and drainage. Nonetheless, at least 400 ha in the deeper parts still retain enough water to attract at least 50,000 waterfowl (Rahmani and Arora 1992). Apart from these two Jheels, there are numerous wetlands beside the canal in an area of about 8,000 ha. To protect them, the Uttar Pradesh government declared a sanctuary of 8,023 ha, including forest land, village land, agricultural fields, and revenue lands. A major part of the Sanctuary (5,300 ha) is private land. The area has been identified as an Important Bird Area due to the presence of globally threatened species such as the Greater Spotted Eagle *Aquila clanga*, *Sarus crane Grus antigone*, and congregations of about 50,000 water birds.

Lakh Bahosi Bird Sanctuary was created in 1988 with the aim of protection and conservation of the wetland with special emphasis to the local and migratory birds, conservation of their natural habitat including aquatic plants and animals. The Bombay Natural History Society, Mumbai (BNHS) has also listed this Sanctuary as one of the 'Important Bird Area' sites. This wetland is

¹Corresponding author

included in the list of wetlands in defined under National Wetland Conservation Programme by the Ministry of Environment & Forest, Govt. of India. Lakh Bahosi Bird Sanctuary consists of two main lakes i.e. Lakh & Bahosi named after respective villages. Approximately 50 thousand water fowls visit this Sanctuary every year between the months of November to March. This Wetland serves as their habitat for nesting as well as breeding for some birds. Thus, this sanctuary has become an attractive destination for the bird lovers as well as tourists since several years. (Figure 1A & 1B)

Efforts to protect bird sanctuaries in Uttar Pradesh include stricter enforcement of conservation laws, promoting sustainable tourism practices, and addressing pollution and habitat loss through government and community initiatives. Protecting bird sanctuaries is crucial for the preservation of biodiversity and the well-being of many bird species. Effective conservation strategies are needed to ensure the survival of these vital habitats. These are several strategies that can help to protect diversity of bird sanctuaries:

Habitat Protection and Restoration: Designating bird sanctuaries as protected areas is one of the most effective ways to safeguard their habitats from human encroachment and exploitation.

Legal Protection, Stronger Enforcement of Laws: Strengthen legal frameworks that protect bird sanctuaries and wildlife, ensuring strict enforcement against illegal activities like poaching, logging, and land conversion.

Community Engagement and Awareness, Local Involvement: Engage local communities in conservation efforts. When communities understand the importance of sanctuaries, they are more likely to protect these areas.

Factors which leads to pollution in Lakh Bahosi Jheel of Bird sanctuary:

1. Agriculture
2. Grazing
3. Fishing
4. Groundwater recharge

Central Lake and Water Source

The Lakh Bahosi Bird Sanctuary is primarily supplied with water from the Ganga Canal. This canal's overflow and seepage accumulate in the two main lakes (Jheels) within the sanctuary, named Lakh and Bahosi. These shallow lakes are vital for the sanctuary's water bird habitat.

The Sanctuary is formed of two oxbow Jheels near the village Bahosi. Both the Jheels, Lakh and Bahosi, are located near the Lower Ganga Canal, so the overflow and seepage of water accumulates in the jheels, resulting in about 600 ha of shallow wetlands perfectly suitable for water birds.




Figure 1A: Location Map of Lakh Bahosi Bird Sanctuary (Source: Google Maps)



Figure 1B: Layout of Lakh Bahosi Bird Sanctuary (Source: Google maps)

Data Collection and Analysis

Tours were conducted to the Lakh Bahosi bird Sanctuary for observation and cataloguing of the flora, in different seasons during 2023 to 2024. Field observations were made on different parts of Sanctuary, photographs of landscapes flora and vegetation types were taken. Healthy and complete plant specimens were cut using scattier and then collected in a polybag and field data on habit and habitat, flowering fruiting time were recorded. For floristic analysis, the plant specimens collected directly from the fields were dried and preserved as reference material by following the ideal herbarium techniques of

Jain and Rao (1977). All specimens were identified with the help of some local flora (Bor 1960, Kanjilal 1966) as well as Duthie's flora and by the help of online websites also. The entire collected voucher specimens were verified through checking with The Plant List (www.theplantlist.org), The International Plant Name Index (IPNI), and Index Kewensis (IK). Specimens were

also matched with the available identified herbarium specimens. In taxonomic treatment, the families are listed as per the Bentham and Hooker's system of classification and all genera within families and species within each Genus are arranged alphabetically (Plate 1 & 2). All results are summarized appropriately and depicted through tables 1 and figures 2.




Plate 1: A- *Cayratia japonica*, B- *Terminalia arjuna*, C- *Physalis peruviana*, D- *Ocimum americanum*, E - *Lantana camara*, F- *Abutilon indicum*




Plate 2: A- *Nymphaea nouchali*, B -*Leersia oryzoides*, C- *Pontederia crassipes*, D- *Saccharum spontaneum*, E- *Alismaplantago-aquatica*, F- *Ipomoea aquatic*

Table 1: List of plant species from the Lakh Bahosi Bird Sanctuary Kannauj, Uttar Pradesh

S.No.	Family Genus and Species	Habit	Habitat	IUCN Status
1	Ranunculaceae <i>Ranunculus sceleratus</i>	H	Sm	LC
2	Malvaceae <i>Abutilon indicum</i> <i>Sida cardifolia</i> <i>Melochia corchorifolia</i> <i>Sida acuta</i> <i>Urena lobata</i>	S S Us H Us	Tr Tr Tr Tr Tr	NE NE LC NE NE
3	Oxalidaceae <i>Oxalis corniculata</i>	H	Tr	NE
4	Meliaceae <i>Azadirachta indica</i>	T	Tr	NE
5	Rhamnaceae <i>Ziziphus oenopila</i> <i>Ziziphus mauritiana</i>	S S	Tr Tr	NE NE
6	Convolvulaceae <i>Evolvulus alsinoides</i> <i>Ipomoea aquatica</i> <i>Ipomea carnea</i>	H H C	Tr Aq Tr/Sm	LC NE NE
7	Cyperaceae <i>Cyperus rotundus</i>	H	Sm/Tr	NE
8	Portulacaceae <i>Portulaca oleracea</i>	H	Tr	LC
9	Verbanaceae <i>Tictona grandis</i> <i>Lantana camara</i>	T S	Tr Tr	EN NE
10	Acanthaceae <i>Peristrophecalyculata</i>	H	Tr	NE
11	Papilionaceae <i>Medicago sativa</i>	H	Tr	NE
12	Primulaceae <i>Anagallis arvensis</i>	H	Tr	NE
13	Lamiaceae <i>Lucas aspera</i> <i>Ocimum americanum</i>	H H	Tr Tr	NE NE
14	Papaveraceae <i>Argemone mexicana</i>	H	Tr	NE
15	Scrophulariaceae <i>Lindenbergia indica</i>	H	Tr	LC
16	Hydrochariataceae <i>Hydrilla verticillata</i> <i>Vallisneria spiralis</i>	H H	Aq Aq	LC LC
17	Ceratophyllaceae <i>Ceratophyllum demersum</i>	H	Sm	LC
18	Myrtaceae <i>Syzygium cumini</i>	T	Tr	NE
19	Mimosaceae <i>Acacia nilotica</i>	T	Tr	LC
20	Phyllanthaceae <i>Phyllanthus reticulatus</i> <i>Phyllanthus amarus</i>	S H	Tr Tr	LC NE

	Vitaceae <i>Cayratia japonica</i>	C	Tr	NE
21	Nymphaeaceae <i>Nymphaea nouchali</i> <i>Nelumbo nucifera</i>	H H	Aq Aq	LC LC
22	Tiliaceae <i>Corchorus trilocularis</i>	H	Tr	NE
23	Zygophyllaceae <i>Tribulus cistoides</i>	H	Tr	NE
24	Pontederiaceae <i>Pontederia crassipes</i>	H	Aq	LC
25	Rubiaceae <i>Oldenlandia corymbosa</i> <i>Anthocephalus chinensis</i>	H T	Tr Tr	LC
26	Fabaceae <i>Dalbergia sissoo</i> <i>Mimosa pudica</i> <i>Cassia tora</i>	T H H	Tr Tr Tr	LC LC LC
27	Mimosaceae <i>Prosopis juliflora</i> <i>Acacia nilotica</i>	T T	Tr Tr	LC LC
28	Combretaceae <i>Terminalia arjuna</i>	T	Tr	LC
29	Cucurbitaceae <i>Coccinia grandis</i> <i>Lagenaria siceraria</i> <i>Cucurbita moschata</i> <i>Luffa cylindrica</i> <i>Luffa echinata</i>	C C C C H	Tr Tr Tr Tr Tr	NE NE NE NE LC
30	Asteraceae <i>Acmella paniculata</i> <i>Ageratina ligustrina</i> <i>Cirsium arvense</i> <i>Eclipta prostrata</i> <i>Cyanthillium cinereum</i> <i>Xanthium strumarium</i> <i>Sonchus asper</i> <i>Eclipta prostrata</i> <i>Parthenium hysterophorus</i> <i>Vernonia cinerea</i> <i>Ageratum conyzoides</i> <i>Tridax procumbens</i> <i>Launaea asplenifolia</i> <i>Erigeron sumatrensis</i> <i>Centipeda minima</i>	H Us H H H H H H H H H H H H H	Tr Tr Tr Tr Tr Tr Tr Tr Tr Tr Tr Tr Tr Tr Sm	LC LC NE LC NE NE NT NE NE NE NE NE NE NE NE
31	Apocynaceae <i>Nerium oleander</i> <i>Thevetia peruviana</i> <i>Alstonia scholaris</i>	S S S	Tr Tr Tr	LC VU NE

	Solanaceae <i>Datura metal</i> <i>Solanum nigrum</i> <i>Physalis peruviana</i> <i>Lycopersicon esculentum</i> <i>Datura stramonium</i>	S H H H H	Tr Tr Tr Tr Tr	NE NE LC LC NE
32	Nyctiganceae <i>Bougainvillea spectabilis</i>	C	Tr	NE
33	Amaranthaceae <i>Achyranthes aspera</i> <i>Alternanthera pungens</i> <i>Alternanthera polygonoides</i> <i>Amaranthus viridis</i> <i>Amaranthus spinosus</i>	H H H H H	Tr Tr Tr Tr Tr	NE LC NE NE NE
34	Chenopodiaceae <i>Chenopodium album</i>	H	Tr	NE
35	Polygonaceae <i>Polygonum plebeium</i>	H	Tr	LC
36	Euphorbiaceae <i>Euphorbia hirta</i> <i>Phyllanthus reticulatus</i> <i>Ricinus communis</i> <i>Croton bonnplandianum</i>	H S S H	Tr Tr Tr Tr	NE LC NE NE
37	Moraceae <i>Artocarpus heterophyllus</i> <i>Ficus religiosa</i> <i>Morus alba</i> <i>Ficus bengalensis</i> <i>Ficus hispida</i>	T T T T T	Tr Tr Tr Tr Tr	NE LC LC NE NE
38	Areceae <i>Phoenix dactylifera</i> <i>Phoenix sylvestris</i> <i>Pistia stratiotes</i>	S S H	Tr Tr Aq	LC LC LC
39	Lemnaceae <i>Lemna perpusilla</i> <i>Spirodela polyrhiza</i>	H H	Aq Aq	NE LC
40	Poaceae <i>Setaria distans</i> <i>Setaria verticillata</i> <i>Saccharum spontaneum</i> <i>Setaria viridis</i> <i>Eclipta alba</i> <i>Cyanodon dactylon</i> <i>Cyperus schoenoides</i>	H H H H H H H	Tr Tr Tr Tr Tr Tr Tr	NE NE NE NE LC NE LC
41	Typhaceae <i>Typha latifolia</i>	H	Aq	LC
42	Aizoaceae <i>Trianthema protulacastrum</i>	H	Tr	NE
43	Molluginaceae <i>Glinus lotoides</i>	H	Tr	LC
44	Salvinaceae <i>Azolla pinnata</i> <i>Azolla caroliniana</i>	H H	Aq Aq	LC LC
45	Marsileaceae <i>Marsilea minuta</i>	H	Aq	LC
46	Pteridaceae <i>Pteris vittata</i>	H	Tr	NE

(H= Herb; C=Climber; S= Shrub; L= Liana, T=Tree; Aq=Aquatic; Tr=Terrestrial; Sm=Semi-aquatic)

RESULTS AND DISCUSSION

Floristic Analysis

A wealthy floristic assemblage with blend of aquatic, wetland and terrestrial plants under different habit ranges of herbs, shrubs, climbers, lianas and trees occur in Lakh Bahosi Bird Sanctuary. There are 110 plant species distributed over 94 genera and 48 families of which 43 families, 84 genera and 97 species are Dicots and 3 families, 7 genera and 9 species are monocots, while the pteridophytes have only 4 species. Evidently, the dicot families representation (87.75%) is almost 4 times that of monocot (6.12%), while the Pteridophytes represent 6.12 %. At Generic level there are 89.36 % dicots, 7.44% monocots and only 6.12% pteridophytes and at species level, 88.18% dicots, 8.18% monocots and 3.63% Pteridophytes. The flora of the sanctuary is richer in

generic composition and diversity as compared to the species diversity among each family. It may therefore be concluded that the small sanctuary possesses a high generic diversity as compared to the family and species with complete dominance of dicots at all levels. (Table 1 and Figure 2)

Habit Types

The maximum representation of the flora is of herbaceous species (72 species) including aquatic herbs, while there are only 14 shrubs, 3 under shrubs, 7 climbers and 14 tree species indicating that the sanctuary flora is predominantly herbaceous with free-floating, suspended amphibious, anchored submerged herbs. Trees are scattered all over contributing to infrequent tree canopy while shrubs and undershrubs mostly border the core water regimes.




Figure 2: Family wise graphical representation of plant species present at Lakh Bahosi Bird Sanctuary

Families, Genera and Species Analysis

Among the five dominant families, Asteraceae was most dominant with 15 species (13.63%) followed by Poaceae with 7 species (6.36%), Solanaceae, Malvaceae and Amaranthaceae with 5 species each having 4.90 % coverage. These together constitute one-third of the sanctuary flora contributing 37 species i.e. 33.63%. The family Asteraceae alone with 15 Genera (15.95%)

suggests a rich representation of ornamental and weeds within the sanctuary comprising grassland and bird food chain resources. The families Poaceae with 5 genera (5.31%) and Malvaceae, Solanaceae, and Amaranthaceae with 5 genera each were also well represented. The most dominant genera with maximum species representation are *Setaria*, *Phyllanthus*, *Ficus bengalensis*, *Azardirachta indica*, *Alstonia scholaris*, *Terminalia arjuna* and *Syzygium jambolanum* etc. (Figure 2)

ACKNOWLEDGEMENTS

The authors are thankful to the Principal Chief Conservator of Forests (PCCF) for providing permission of field visit of the protected sites of Lakh Bahosi and Head, Department of Botany, University of Lucknow for laboratory facilities.

REFERENCES

- Agnihotri P., Hussain T. and Maliya D. 2008. Aquatic flora of some notified Bird Sanctuaries of Uttar Pradesh. *Indian Forester*, **134**: 1398-1401.
- Duthie J.F., 1903-1915. Flora of the upper Gangetic plain, and of the adjacent Siwalik and sub Himalayan tracts (Vol. **1-3**). Superintendent of Government Printing.
- IUCN, 2012. Guidelines for application of IUCN red list criteria at regional and national levels: version 4.0.
- Garg A. and Singh P., 2018. Botanical Survey of India. *Indian Journal of Forestry Floristic Diversity of Nawabganj Bird Sanctuary*.
- Jain S.K. and Rao R.R., 1977. *A Handbook of Field and Herbarium Methods*. Today & Tomorrow's Printers and Publishers, New Delhi.
- Joshi P., 2014. Study on Waders and Wetland Bird Diversity, and their Habitat Selection of Some the Fresh Water Resources around Yavatmal City, Maharashtra, India. *Nature & Environment*, **19**(2): 188-194.
- Kanaujia A., Kumar A., Kushwaha S. and Kumar A., 2015. Diversity of Waterbirds in Lucknow District, Uttar Pradesh, India. *International Journal of Science and Research (IJSR)*, **4**(1): 862-866.
- Koladiya M.H., Mahato A.K.R. and Pardeshi M. 2014. Status of Avi-fauna in Lakhara Wetland, Kachchh Biosphere Reserve, Gujarat, India. *Cibtech Journal of Zoology*, **3**(1): 37-41.
- Kanjilal P.C., 1996. A forest flora for the Plants of Uttar Pradesh (Part II & III) Lucknow, U.P., India.
- Kumar D.R. and Kanaujia A., 2015. Lakh Bahosi Bird Sanctuary: A Waterbird Abode.
- Kumar A. and Sati J.P., 2003. Check list of India wetland in Envis News letter Avian Ecology & Inland wetland Buceros., **8**(1): 1-29.
- Singh R.I. and Tomar D.S., 2010. Nawabganj Pakshi Vihar ki Prabandh Yojna 2000-2010 Vanya Jeev Parikshan Sangathan U.P. Lucknow
- Sinha B.K., 2005: Floristic diversity and vegetation composition of Lakh-Bahosi wetlands of Kannauj, Uttar Pradesh.