



## PHYTOCHEMICAL INVESTIGATION AND ANTIBACTERIAL ACTIVITY OF LEAF EXTRACT OF CASSIA ALATA LINN MEDICINAL PLANT

RAHUL VERMA<sup>a1</sup> AND SHAILENDRA SINGH<sup>b</sup>

<sup>ab</sup>Department of Chemistry, T.D.P.G College, Jaunpur, (U.P) India

### ABSTRACT

The Leaves of *Cassia alata* Linn were successively extracted with n-hexane and ethyl acetate using the soxhlet method. Preliminary phytochemical screening of the extracts revealed the presence of free anthraquinones, flavonoids, steroids and saponins. The n-hexane crude extract exhibited some anti-bacterial activity against *Yersinia enterocolitica*, *Streptococcus pneumonia* and *Salmonella typhi*. Anti-fungal activities against *Microsporum audouini* and *Trichophyton mentagrophytes* were also exhibited. A synergic test of n-hexane and ethyl acetate extracts showed an improved sensitivity against *Shigella sonnei* and *Strep pneumonia*. A confirmatory phytochemical analysis performed on the most mobile tlc isolate ( $R_f$  0.94) from the n-hexane extract revealed the presence of steroidal saponin. This was found to be active against strep pneumonia.

**KEYWORDS:** *Cassia alata*, Phytochemistry, Steroids and Antimicrobial Activity

A number of plant have been reported in literature to possess antibacterial and antifungal properties. Among the plants with such activities is *Cassia alata* Linn. It is one of the 18,000 species of leguminaceae family. This plant is a small perennial shrub, which grows rapidly in full sun on a wide range of soil from a height of 3m to 4.5m with a straight, wooden stem (Giron, 1991). The large leaves are bilateral – symmetrical opposed and fold together at night. The fruit is a pod with a woody brown shell about 120cm long and hard to touch *Cassia alata* is common plant in Japan, Indonesia, Mexico, Ghana, Trinidad and Nigeria (Giron, 1991). The Leaves have been reported to be sudorific, diuretic and purgative. According to Bhat, Eterjere, and Okdipo (1990), the leaves have been used for the treatment of constipation, ringworm and other skin diseases. The leaves in decoction are also used to treat bronchitis and asthma. The use of the leaves of the plant as anti-venom and as an abortifacient are also known (Bhat *et al.*, 1990)

The present work was therefore designed to do the following:

- i - Identify the phytochemicals present in the non-polar n-hexane and ethyl acetate extracts of the leaves of *Cassia alata* Linn.
- ii – Confirm the antibacterial and antifungal activity of the non – polar extracts of the leaves of the plant.
- iii – Purify the crude extracts in order to identify the principles responsible for the activity.

### EXPERIMENTAL

#### Collection and Preparation of Plant Materials

The fresh leaves of the plant *Cassia alata* Linn, were collected from the rural area of Jaunpur district (UP) and identified by Taxonomist of Botany Department T.D.P.G. College Jaunpur, and herbarium was deposited as assigned that voucher specimen no. 126.

#### Extraction

200g of the powdered leaves of *Cassia alata* Linn was successively extracted with soxhlet extractor using each of the following solvents: n-hexane (30-40°C) and ethyl acetate (30-40°C) in that order. The extraction was carried out at near the boiling point of the extracting solvent. Each extract was concentrated in – vacuo with the aid of a rotary evaporator. The respective extract concentrates were kept in desiccators to dry for at least 3 days before further tests were carried out on them.

#### Preliminary Phytochemical Screening

Preliminary phytochemical screening was carried out on the crude n-hexane and ethyl acetate extracts. Known standard procedures for phytochemical screening were adopted to test for the presence of cardiac glycosides, glycosides, saponins, anthraquinones, tannis, terpenes, steroids, alkaloids and flavanoides (sofo-wora, 1993) .These same phytochemicals were tested for in the tlc fraction obtained from the n-hexane extract.

### Chromatographic Purification of n-Hexane Crude Extract

To purify the n-hexane extract, various fractions were spotted on a silica tlc plates and developed using various solvent mixture. It was observed that a 17:3 chloroform n- hexane mixture gave the best resolution. This solvent mixture was used to resolve these fractions. The most mobile yellow band ( $R_f$  0.94) on a preparative tlc (Braith-Waite and Smith, 1985) plate was isolated with hexane as a yellow oil. This was further dried several days in a desiccator.

### Antimicrobial Studies

Pure clinical isolates of streptococcus pneumonia, salmonella typhi, E.coli, Yersinia enterocolitica, shigella sonnei, microsporum audouinii and trichophyton mentagrophyte were obtained from the National Botanical Research Institute, (NBRI). The bacteria were grown on nutrient agar slants in an incubator at 37°C for 24 hours while the fungi were sub-cultured on the slant sabourand dextrose agar and incubated at 25°C for 48 hours. Growth of micro-organism in the broth was indicated by turbidity. The turbidity produced was adjusted to match 0.5 Mc for land standards ( $10^8$  cfu /ml), which was further adjusted to ( $10^5$  cfu /ml) for bacteria and fungi respectively.

The cork and bore diffusion method was used to carry out the antimicrobial screening. Solutions of the crude extracts were prepared for n-hexane and ethyl acetate extracts. 20mg /ml of each was prepared.

Mean while, inoculation of the prepared plates with the organism was done using wire- loop to transfer strands of the organism into the plates followed by a cross streaking with the same wire loop to achieve uniform spread of the microbes on the plates. These were done for both the nutrient agar (NA) and Sabourand dextrose agar (SDA) plates. After about 30 min, wells were punched on the plates using a sterile cork borer of 5mm diameter.

A 0.1 ml of the crude extract solution was introduced into each appropriately labeled well. Tetracycline of the same concentration as the extracts was used as control for NA plate containing bacteria while climaxol was used as control for SDA plates containing the fungi. After proper diffusion had taken place the plates were incubated at 37°C for 24 hours for bacteria and 48 hours at 25°C for fungi. Zones of inhibition of growth were determined by linear measurement of diameter.

This same procedure was adopted for the pure yellow oil fraction obtained from the preparative tlc studies.

### Test for Synergism

Synergic effect of the extracts of the leaves of a *Cassia alata* Linn was tested on the human pathogens used for antimicrobial activity test. This was carried out by measuring equal amount of the n-hexane and ethyl acetate extract and thus preparing a solution of the mixture. The cork and bore diffusion method was repeated for the mixture to test its sensitivity.

### RESULTS AND DISCUSSION

Table 1 shows the phytochemical constituents of the leaves of *Cassia alata* Linn. Anthraquinones, saponins, steroids and flavonoids were present in both solvents used for extraction. On the other hand alkaloids, tannins, terpene, glycosides and cardiac glycosides were not detected in the solvents.

Anthraquinones according to Mohammad (1994), are known for their purgative (cathartic) property, especially in habitual constipation. Saponins on the other hand are known to possess both antimicrobial and anti-inflammatory properties (Margineanu *et al.*, 1976 ; Singh *et al.*, 1974). They are also known to serve as precursor of steroidal substance with a wide range of physiological activities. Pharmacologically, some flavonoids have also been found to exhibit anti-tumor, anti-bacterial and antifungal, properties (Trease and Evans, 2002; Finar, 1980). The presence of these phytochemicals may account for the spectrum of activity of the plant.

The n-hexane extract of the leaves of *Cassia alata* Linn exhibited some antibacterial activity against yersinni enterocolitica streptococcus pneumonia and Salmonella typhi.

Also ethyl acetate extract demonstrated antibacterial activity against Strep. Pneumonia, yersinnia enterocolitica and E.coli. Both n- hexane and ethyl acetate extracts were found to be active against *Microsporum audouinii* and *Trichophyton mentagrophyta*. These were the two used for the test (Table 2). A synergic test of n-hexane and ethylacetate crude extracts show an improved against.

*Shigella sonnei* and *Streptococcus pneumonia* (Table 2). Preparatory tlc carried out on the n- hexane extract resulted in the isolation of a yellow and oily component ( $R_f$  0.94): a confirmatory phytochemical analysis performed on this isolated revealed the presence of steroidal saponin. This was found to be active against strep. Pneumonia (Table 3).

The results of the antimicrobial activity of the extracts are in agreement with the uses of the extracts of the leaves of *Cassia alata* Linn in traditional medicine for

the treatment of constipation, sexually transmitted disease (STD), bronchitis and asthma, ring worm and other skin disease. These results further substantiate the literature reports that extracts of the leaves of *Cassia alata* Linn have both antibacterial and anti-fungal properties (Okafor

*et al.*, 2001; Ibrahim and Osman, 1995; Somchit *et al.*, 2003). The leaves of the plant therefore appear to be a potential source of broad spectrum antibiotics.

**Table 1: Phytochemical constituents of the leaf of *Cassia alata* Linn**

Metabolites Extracts	1	2	3	4	5	6	7	8	9
HE	-	++	-	++	++	-	-	-	++
EAE	-	++	-	++	++	-	-	-	++

HE = n-hexane extract, EAE = ethylacetate extract

- |               |                        |               |
|---------------|------------------------|---------------|
| 1. Alkaloids  | 2. Free anthraquinones | 3. Tannins    |
| 4. Saponins   | 5. Steroids            | 6. Terpenes   |
| 7. Glycosides | 8. Cardiac glycosides  | 9. Flavonoids |

**Table 2: Results of antimicrobial activity test of Crude extracts from the leaf of *Cassia alata* Linn.**

Test Organism	Diameter of Zone Inhibition (mm)				
	HE	EAE	HE/EAE	TCN	C
<i>Strep pneumonia</i>	20.0	10.0	22.0	13.0	NA
<i>Shigella sonnei</i>	6.0	8.0	20.0	13.0	NA
<i>Yersinia enterocolitica</i>	12.0	10.0	12.0	11.0	NA
<i>Escherichia coli</i>	7.0	10.0	7.0	12.0	NA
<i>Salmonella typhi</i>	12.0	9.0	6.0	15.0	NA
<i>Microsporum audouinii</i>	25.0	22.0	16.0	NA	18.0
<i>Trichophyton mentagrophyte</i>	23.0	22.0	23.0	NA	20.0

HE= n-hexane extract, EAE = Ethyl acetate extract.

HE/ EAE = a mixture of equal amount of HE and EAE

TCN = Tetracycline (control against bacteria )

C = Climaxol (control against fungi)

NA = Not applicable

**Table 3: Sensitivity test of the pure component of n- hexane extract of the leaf of *Cassia alata* Linn.**

Zones of Inhibition (mm)			
Test organism	HE	TCN	C
<i>Strep pneumonia</i>	25.0	17.0	NA
<i>Yersinia enterocolitica</i>	20.0	30.0	NA
<i>Salmonella typhi</i>	0.0	17.0	NA
<i>Microsporum audouinii</i>	29.0	NA	38.0
<i>Trichophyton mentagrophyte</i>	10.0	NA	17.0

HEf = Preparatory tlc fraction from n-hexane extract.

## CONCLUSION

In conclusion, the antimicrobial screening gave some justification for the use of the leaves of *Cassia alata* Linn in the traditional medicine. The results have also shown that n-hexane and ethyl acetate are good solvents for the extraction of phytochemicals from the leaves of *Cassia alata* Linn. Spectroscopic studies are in progress to characterize the active principles in the leaves of the plant.

## ACKNOWLEDGMENTS

The authors are thankful to Director N.B.R.I Lucknow, U.P. for providing spectral analysis of the compound. We are thankful to Head, Department of Chemistry T.D.P.G. College, Jaunpur (U.P.) for providing necessary facilities to complete the work.

## REFERENCES

- Giron L.M., 1991. Ethno-botanical survey of the medicinal flora used by the Guatemalan. *Journal Ethnopharmacology*, **34** (2-3): 173-187.

- Farnsworth N.R and Bunyaphatsara, 1992. Thai medicinal plants, Recommended for Primary health care system, Prachachon, Bangkok.
- Bhat R.B, Eterjere E.O. and Oladipo V.T., 1990. Ethnobotanical studies from central Nigeria econ. Bot. **44**(3): 382-390.
- Braithwaite A. and Smith F.J., 1985. Chromatographic Methods (4<sup>th</sup>Ed), New York, Chapman and Hall Ltd., pp 140-150.
- Sofowora A., 1993. Medicinal Plants and Traditional medicine in Africa. Ibadan, John Wiley and Sons Ltd. P 209.
- Trease G.E and Evans W.C., 2002. Pharmacognosy (5<sup>th</sup>ed). London W.B Saunders.
- Mohammed, 1994. Text book of pharmacognosy. CBS publishers & distributors, 4596 / 1A, 11-Daryaganji, New Delhi; 110002, P.107.
- Margineanu V.C., Cucu V., Grecu L. and Parvu C., 1976. Nig J. Bot., **19**(1): 156-162.
- Singh S., Ehana N.M. and Dhar M.M., 1974. Nig J. Bot: **19** (1):156-162.
- Finar I.L., 1980. Chemistry- stereochemistry and chemistry of natural products. (5<sup>th</sup>ed ). London. Longman group Ltd. P. 877.
- Okafor J.I., Eze E.A. and Njoku S.O.U., 2001. Antifungal activities of the leaves of baphia, Nitida, *Cassia alata*, ficus exasperate and gossypium arboreum. Nig. Journal of Nat. Product and Med., **5**:59-60.
- Ibrahim D. and Osman H., 1995. Antimicrobial activity of *Cassia alata* from Malaysia, Journal of Ethnopharmacology, **45**(3): 151-156.
- Somchit M.N., Reezal I., Elysha N.I. and Mutalib A.R., 2003. Invitroantimicrobial activity of ethanol and water extracts of *Cassia alata*. Journal of Ethnopharmacology, **84**(1):1-4.